



Reptiles of the King Abdulaziz Royal Nature Reserve, east-central Saudi Arabia: insights and conservation implications

¹Ryan van Huyssteen, ¹Melissa A. Petford, ^{1,2}Marius Burger, ^{1,3,4}Jiří Šmíd, ⁵Abdulrahman S. Alzahrani, ^{5,*}Abdullah M. Alowaifeer, ¹Phoebe Mottram, and ¹Jerome Y. Gaugris

¹FLORA FAUNA & MAN, Ecological Services Ltd, Tortola, BRITISH VIRGIN ISLANDS ²African Amphibian Conservation Research Group, Unit for Environmental Sciences and Management, North-West University, Potchefstroom, SOUTH AFRICA ³Department of Zoology, National Museum, Prague, CZECH REPUBLIC ⁴Department of Zoology, Faculty of Science, Charles University, Prague, CZECH REPUBLIC ⁵Environmental and Sustainability Department, King Abdulaziz Royal Nature Reserve Development Authority, Riyadh, SAUDI ARABIA

Abstract.—The distribution of herpetofauna on the Arabian Peninsula is generally poorly known, particularly in Saudi Arabia. The King Abdulaziz Royal Nature Reserve (KARNR) in east-central Saudi Arabia is in an area predicted to have high reptile species richness, yet there is no baseline reptile checklist for the reserve. Knowing which species occur within a protected area is vital for ensuring that conservation strategies and long-term monitoring are effective. Here, we provide the first detailed list of reptile fauna occurring in the KARNR. This study recorded a total of 31 species from 1,551 observations, including 25 lizards and six snakes. A desktop study identified three additional species, bringing the total number of herpetofaunal species known in the KARNR to 34. Two of the species found in the reserve are of conservation concern: *Uromastyx aegyptia* (VU) and *Tropicolotes wolfgangboehmei* (DD). Through species accumulation curves we were able to determine that the sampling methods implemented during the survey were adequate, and we predicted that the species richness of the reserve is likely as much as 38 species. Prior to this study, only 82 reptile records (30 species) were known from the study area, so we add a significant number of new records to the reserve and an additional four species that were previously unknown from the region.

Keywords. Arabian Peninsula, baseline, iNEXT, protected area, species accumulation curves, species richness, Squamata, *Tropicolotes wolfgangboehmei*, *Uromastyx aegyptia*

Citation: van Huyssteen R, Petford MA, Burger M, Šmíd J, Alzahrani AS, Alowaifeer AM, Mottram P, Gaugris JY. 2024. Reptiles of the King Abdulaziz Royal Nature Reserve, east-central Saudi Arabia: insights and conservation implications. *Amphibian & Reptile Conservation* 18(1&2): 91–106 (e335).

Copyright: van Huyssteen, et al. 2024. This is an open access article distributed under the terms of the Creative Commons Attribution License [Attribution 4.0 International (CC BY 4.0): <https://creativecommons.org/licenses/by/4.0/>], which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. The official and authorized publication credit sources, which will be duly enforced, are as follows: official journal title *Amphibian & Reptile Conservation*; official journal website: amphibian-reptile-conservation.org.

Accepted: 2 September 2024; **Published:** 25 November 2024

Introduction

Reptile species richness on the Arabian Peninsula is highest in the mountainous regions along the south-western, southern, and eastern parts of the peninsula (Carranza et al. 2021; Šmíd et al. 2021). However, the central desert regions also have relatively high species richness and phylogenetic diversity (Cox et al. 2012; Šmíd et al. 2021; Tamar et al. 2023). The Kingdom of Saudi Arabia, which covers most of the Arabian Peninsula, is well known for its large expansive deserts and a number of regional inventories of reptiles have been compiled over the past few decades (i.e., Al Sadoon et al. 2016, 2017; Aloufi et al. 2021, 2022; Alshammari and Busais 2020; Alshammari et al. 2017; Alshammari and Ibrahim 2015; Aloufi and Amr 2015; Cunningham 2010; Farag and Banaja 1980; Tilbury 1998). Despite these inventories, large areas of Saudi Arabia are poorly

surveyed for reptiles at the landscape scale, so our current knowledge of the distribution and community composition of Saudi Arabia's reptiles is inadequate (Šmíd et al. 2021).

The King Abdulaziz Royal Nature Reserve (KARNR) in east-central Saudi Arabia was established in 2018 and is one of seven royal reserves in the country (<https://www.karnr.gov.sa/>). The reserve covers a heterogenous landscape of just over 28,000 km², and this region is predicted to have high terrestrial reptile species richness (Alatawi et al. 2020; Ficetola et al. 2013; Šmíd et al. 2021). However, there have been no dedicated surveys on the reptile species richness of the area. The lack of such inventories in protected areas is detrimental to reptile conservation as there are no baseline data to inform conservation strategies and long-term monitoring methods in the reserve (Margules and Pressey 2000; Mihoub 2017).

Correspondence. *rv.huyssteen@florafaunaman.com (RVH); a.alowaifeer@karnr.gov.sa (AMA)

Reptiles of the King Abdulaziz Royal Nature Reserve

This study aimed to provide an adequate baseline of reptile species richness in the KARNR. To achieve this, we conducted field work on five occasions between November 2022 and October 2023 for a total of 46 days, which equated to 139 person-days in the field. A desktop study was also conducted to identify any reptile species previously recorded from the area. We tested the adequacy of our survey effort using species accumulation curves and estimated diversity indices.

Materials and Methods

Study Area

The KARNR is situated in east-central Saudi Arabia. It is located between the northern Riyadh region and northwestern Eastern Province and covers an area of just over 28,000 km² (Fig. 1). Our study area was the KARNR plus a 10 km buffer around the boundary of the reserve. According to the Köppen-Geiger classification, this region is categorized as a hot desert (Peel et al. 2007). The major geomorphological features of the study area are the Al Summan plateau in the north of the reserve, the Ad Dahna sand desert that crosses the reserve nearly transversely in the middle, and the Tuwaiq Escarpment, a mountain ridge that cuts through the southern part of the reserve (Almalki et al. 2022; Vincent 2008). The influence of these sandy and rocky formations has resulted in a high habitat heterogeneity for the region, with rocky areas (such as escarpments and rock outcrops), open plains (stoney and gravel plains), dunes (mobile and vegetated dunes), and wetland areas and drainage lines (wadis and rawdhats) (Fig. 2). A general north/south moisture gradient is evident from the vegetation and plant communities in the study area, with



Fig. 2. Several landscape features of the King Abdulaziz Royal Nature Reserve, showing some of the habitat diversity in the region.

the southern areas having more trees and rawdhat areas (Ghazanfar and Fisher 1998).

Methods

Field surveys. Three primary field surveys were carried out for 21 days from 23 November to 13 December 2022, 18 days from 3 to 20 March 2023, and eight days from 28




Fig. 1. The King Abdulaziz Royal Nature Reserve study area and its location in the Arabian Peninsula. Blue dots show sampling points visited during the 2022/23 survey, and white stars show reptile trap array locations.

September to 5 October 2023. Two opportunistic surveys were also conducted for a five-day period, which formed part of larger scale rapid biodiversity overviews and were more limited in effort than the three primary surveys. One was from 17 to 21 February 2023 and another was from 8 to 12 June 2023. Collectively, these covered a total of 57 survey-days and 139 person-days. Sampling methods during the field surveys included active searching (both diurnal and nocturnal, see below) and trapping (6 March until 19 March 2023). Observations recorded by team members surveying other taxa were also included as supplementary records. The February 2023 opportunistic survey also targeted wetland areas after a good rain season to check for the occurrence of amphibians in the study area. The June 2023 opportunistic survey was conducted during a hot period of the year and had a specific focus on nocturnal active searches. Reptiles were recorded to the species level when possible. Occasionally some individuals could only be identified to genus when species assignment was uncertain. When feasible, DNA barcoding was conducted to aid in the identification in these instances. For each record, a GPS location (with an average accuracy of 5 m) was taken using Garmin Montana i750 GPS units, and specimen abundance was noted.

Active searching was conducted at 374 specific point locations across the reserve (Fig. 1). These points were selected by overlaying a 1 km sampling grid across the study area and using a multistage, stratified, semi-random sampling approach. The northern section of the reserve, Al Summan, was only surveyed during November–December 2022 and September 2023 due to logistical constraints. The southern section of the reserve was surveyed during all sampling seasons. Diurnal surveys were conducted between 0600 and 1900 h, while nocturnal surveys took place between 1900 and 2300 h.

Eight trap arrays were deployed within the study area. The placement of trap arrays was limited to areas near main roads due to logistical convenience. Each trap array consisted of three 10 m drift fences of robust plastic, arranged in a linear design as described by Mendes et al. (2015), together with four 20 L buckets sunk into the ground as pitfall traps, and six wire-mesh funnel traps positioned on each side of the drift fence between the pitfall traps (Fig. 3). The funnel traps were covered with shade netting to provide shelter for captured animals, while moistened leaf-litter and cloth were added to the buckets to provide shelter and moisture in the pitfall traps. These traps were checked daily for captured animals. The eight trap arrays were installed for 13 nights, equating to 104 trap array-nights.

A desktop survey was also conducted to consolidate historical reptile records for the study area. This was achieved by consulting various literature sources (e.g., Aloufi et al. 2019; Al-Sadoon 1988; Al-Sadoon 1989; Gasperetti 1988; Kordges 1998; Salvador 1982; Šmíd et al. 2021) and online databases (e.g., Global Biodiversity




Fig. 3. (A) An example of a trap array showing drift fence and pitfall trap. (B) Detail of a funnel trap with shade netting cover.

Information Facility at <http://GBIF.org> and iNaturalist at <http://inaturalist.org>. All photographic records obtained during the 2022/23 survey were uploaded to iNaturalist and these are available in Supplemental Table S1.

DNA sequencing. Some species of Arabian reptiles are challenging to identify based on morphological characters alone. We therefore sequenced a short fragment of one mitochondrial gene and applied DNA barcoding methods to verify the identification in the field. Of the species recorded during the surveys, we DNA barcoded members of the following genera: *Acanthodactylus* and *Mesalina* (both Lacertidae); *Pseudotrapelus* and *Trapelus* (both Agamidae), *Hemidactylus* and *Stenodactylus* (both Gekkonidae), *Scincus* (Scincidae), and *Ptyodactylus* (Phyllodactylidae). Tissue samples were obtained by taking a small piece of tail. All samples were preserved in 96% ethanol and DNA was extracted using the Qiagen DNeasy Blood and Tissue Kit (Qiagen, Valencia, California, USA) following the manufacturer's protocol. We PCR-amplified and sequenced the ribosomal 12S rRNA (12S) region of the mitochondrial DNA for all groups except the agamids, for which we sequenced the 16S rRNA (16S). Primers and PCR conditions are detailed in Šmíd et al. (2021). The PCR products were sequenced from both directions at Macrogen (Amsterdam, the Netherlands), and raw sequence data were edited and assembled in Geneious v.11 (Kearse et al. 2012). The identification of each specimen was confirmed using the BLAST algorithm of the National Center for Biotechnology Information website (<https://blast.ncbi.nlm.nih.gov/>).

Species Accumulation Curves

Total survey. To assess our survey effort in the KARNR, we used species accumulation curves calculated in R (R




Fig. 4. Records of reptiles found during the 2022/23 survey period (purple circles) and from the desktop study (blue circles) in the King Abdulaziz Royal Nature Reserve.

Core Team 2022) using the R studio console (RStudio team 2022) with the iNEXT package (Hsieh et al. 2016). Species accumulation curves were plotted and extrapolated to double the original survey effort (Hsieh et al. 2016), and the order of the observations made during the survey were randomized 100 times in a bootstrap fashion. The estimated species richness was calculated in iNEXT along with the standard error and lower and upper confidence levels. Shannon and Simpson diversity indices were also calculated. Four broad-scale reptile habitat types were identified for the KARNR: (1) Dunes and dune valleys, (2) Hills, outcrops, and escarpments, (3) Plains, and (4) Wadis, rawhdats, and pans. These habitat types were defined based on vegetation and topographic characteristics. For each habitat type, predictive species accumulation curves were provided to assess the sampling effort for each of the broad habitat types.

The survey species richness was then evaluated in terms of completeness according to the iNEXT estimated species richness. A percentage of the completeness score was calculated as $\text{SurveySR}/\text{iNEXT Estimated SR} \times 100$. Any result greater than 90% is regarded as “close to completely surveyed,” since it can be unlikely that all species could be recorded during a finite survey (Chao and Jost 2012; Moreno and Halfiter 2001; Thompson et al. 2007). Note that the value calculated is an estimate based on the species accumulation curve, and therefore dependent on the sampling effort.

Survey effort was also assessed for each habitat type to evaluate diversity at the broad-scale habitat level. For this, the vegetation class for each species record was extracted from a vegetation layer generated for the suite of studies conducted for the present project (Van Rooyen et al. 2023) using QGIS 3.22.8 (QGIS Development Team 2023). Each vegetation type class represented in

the reserve was then grouped into broad habitat units of Dunes and dune valleys; Plains; Outcrops, hills and escarpments; and Wadis, rawhdats and pans. The methods above were used to create species accumulation curves and estimate species richness.

Results

A total of 1,551 reptile observations of 31 species were made during the 2022/2023 survey period, composed of 25 lizards and six snakes (Table 1, Supplemental Table S1, Figs. 4–9). Four species had not been previously recorded in the area: *Trapelus agnetae*, *Hemidactylus granosus*, *Mesalina saudiarabica*, and *Diplometopon zarudnyi*. The desktop study of published and online sources produced 82 records of 30 species, three of which were not recorded during our survey: *Platyceps rhodorachis*, *Myriopholis macrorhyncha*, and *Walterinnesia morgani*. Therefore, the total known reptile species richness for the KARNR is currently 34 species.

Most reptile observations were recorded via active searching ($n = 1,400$). Fifty-three individuals were captured by trap arrays, with the most common being *Acanthodactylus ophiodurus* ($n = 11$). The remaining 91 records were all incidental. The most common species encountered was *Bunopus tuberculatus* ($n = 579$), followed by *Uromastix aegyptia* ($n = 367$). Two species were recorded only once during the survey: *Acanthodactylus hardyi* and *Echis coloratus* (Table 1).

Two species of conservation concern were found within the reserve borders: *U. aegyptia* (Vulnerable) and *Tropicolotes wolfgangboehmei* (Gekkonidae; Data Deficient) (IUCN 2023). Although *U. aegyptia* was very common throughout the reserve, only two *T. wolfgangboehmei* were found. One of the localities for *T.*

Table 1. Checklist and abundance of reptile species recorded in the King AbdulAziz Royal Nature Reserve. The International Union for Conservation of Nature (IUCN 2023) listings of conservation status refer to the following categories: Not Evaluated (NE), Least Concern (LC), Data Deficient (DD), Vulnerable (VU), and Endangered (EN).

Scientific name	Common name	IUCN status	Number of observations	Recorded during survey	Recorded prior to survey
SAURIA		Lizards			
Family Agamidae					
<i>Phrynocephalus arabicus</i>	Arabian Toad-headed Agama	LC	26	Y	Y
<i>Pseudotrapelus tuwaiqensis</i>	Tuwaiq Agama	NE	2	Y	Y
<i>Trapelus agnetae</i>	Northern Arabian Plain Agama	LC	4	Y	N
<i>Trapelus ruderatus</i>	Olivier's Agama	LC	4	Y	Y
<i>Uromastix aegyptia</i>	Egyptian Spiny-tailed Lizard	VU	367	Y	Y
Family Gekkonidae					
<i>Bunopus tuberculatus</i>	Baluch Ground Gecko	LC	579	Y	Y
<i>Cyrtopodion scabrum</i>	Rough-tailed Gecko	LC	4	Y	Y
<i>Hemidactylus granosus</i>	West Arabian Half-toed Gecko	NE	5	Y	N
<i>Stenodactylus doriae</i>	Middle Eastern Short-fingered Gecko	LC	44	Y	Y
<i>Stenodactylus slevini</i>	Slevin's Short-fingered Gecko	LC	16	Y	Y
<i>Trigonodactylus arabicus</i>	Arabian Web-footed Sand Gecko	LC	13	Y	Y
<i>Tropiocolotes wolfgangboehmei</i>	Wolfgang's Sand Gecko	DD	2	Y	Y
Family Phyllodactylidae					
<i>Ptyodactylus cf. hasselquistii</i>	Hasselquist's Fan-footed Gecko	LC	107	Y	Y
Family Lacertidae					
<i>Acanthodactylus boskianus</i>	Bosc's Fringe-fingered Lizard	LC	24	Y	Y
<i>Acanthodactylus hardyi</i>	Hardy's Fringe-fingered Lizard	NE	1	Y	Y
<i>Acanthodactylus ophiodurus</i>	Snake-tailed Fringe-fingered Lizard	LC	54	Y	Y
<i>Acanthodactylus schmidti</i>	Schmidt's Fringe-fingered Lizard	LC	57	Y	Y
<i>Mesalina brevirostris</i>	Short-nosed Desert Lizard	LC	102	Y	Y
<i>Mesalina</i> lineage 4	Small Spotted Desert Racer	-	46	Y	Y
<i>Mesalina saudiarabica</i>	Arabian Short-nosed Desert Lizard	NE	3	Y	N
Family Scincidae					
<i>Chalcides ocellatus</i>	Ocellated Skink	LC	8	Y	Y
<i>Scincus mitranus</i>	Arabian Sandfish	LC	12	Y	Y
<i>Scincus conirostris</i>	Sandfish Skink	NE	12	Y	Y
Family Trogonophidae					
<i>Diplometopon zarudnyi</i>	Zarudny's Worm Lizard	LC	23	Y	N
Family Varanidae					
<i>Varanus griseus</i>	Desert Monitor	LC	7	Y	Y

Reptiles of the King Abdulaziz Royal Nature Reserve

SERPENTES	Snakes				
Family Boidae					
<i>Eryx jayakari</i>	Arabian Sand Boa	LC	2	Y	Y
Family Colubridae					
<i>Platyceps rhodorachis</i>	Braid Snake	LC	0	N	Y
<i>Spalerosophis diadema</i>	Diadem Snake	LC	5	Y	Y
Family Elapidae					
<i>Walterinnesia morgani</i>	Black Desert Cobra	LC	0	N	Y
Family Leptotyphlopidae					
<i>Myriopholis macrorhyncha</i>	Beaked Thread Snake	LC	0	N	Y
Family Psammophiidae					
<i>Malpolon moilensis</i>	Hooded Malpolon	LC	6	Y	Y
<i>Psammophis schokari</i>	Afro-Asian Sand Snake	LC	4	Y	Y
Family Viperidae					
<i>Cerastes gasperettii</i>	Arabian Horned Viper	LC	11	Y	Y
<i>Echis coloratus</i>	Palestine Saw-scaled Viper	LC	1	Y	Y
Total observations and species counts			1,551	31	30
Total species for KARNR				34	

wolfgangboehmei was inside the 10 km buffer zone from a known site, whereas the other was inside the reserve and represents a new locality record for this species.

DNA Barcoding

We DNA barcoded a total of 95 samples in the families Agamidae (four samples), Gekkonidae (13 samples), Lacertidae (59 samples), Phyllodactylidae (seven samples), and Scincidae (12 samples). The results mostly confirmed our field identifications. However, two species were misidentified in the field and were only revealed by the DNA analysis. These were one specimen of *Acanthodactylus hardyi* and three specimens of *Mesalina saudiarabica*, both from the vicinity of the Ad Dahna sand stretch. It is important to note the majority of *Mesalina brevirostris* specimens were not DNA barcoded, so some of these may be incorrectly identified as *M. saudiarabica*. Details on the DNA barcoding results along with GenBank accessions of the newly sequenced samples are given in Supplemental Table S2.

Species Accumulation Curves

The predictive species accumulation curve indicates that the sampling methods employed during the field surveys were adequate, with the curve reaching an asymptote (Fig. 10) and an estimated 97.89% of reptile species

being recorded, based on the iNEXT completeness score (Table 2). The species accumulation curve predicts that the species richness for the KARNR is between 31 and 38 species (LCL = 31; UCL = 37.84; Table 2).

Dunes and dune valleys are the least well sampled of the broad habitat units, with the accumulation curve not reaching an asymptote (Fig. 11) and an estimated 63.75% of the reptile species being recorded based on the iNEXT completeness score (Table 2). Plains are relatively well sampled, but the iNEXT completeness score of 88.75% suggests that additional species are likely to be found in this area, with an estimated species richness between 21 and 38 (LCL = 21.0; UCL = 38.05). Wadis, rawhdats, and pans also appear to be well sampled with an iNEXT completeness of 94.93%. Hills, outcrops, and escarpments are the most well sampled habitat type with iNEXT completeness scores of over 98% and accumulation curves approaching asymptotes (Table 2; Fig. 11).

Discussion

There are now 34 species of reptiles confirmed to occur within the KARNR. Thirty-one of these species were found during our survey, four of which had not previously been recorded in this area. This study contributes a substantial number of observations for the region, with 1,550 individual records across the reserve, compared




Fig. 5. (A) *Phrynocephalus arabicus*, (B) *Pseudotrapelus tuwaiqensis*, (C) *Trapelus agnetae*, (D) *Trapelus ruderatus*, (E) *Uromastyx aegyptia*, (F) *Uromastyx aegyptia* burrow, (G) *Uromastyx aegyptia* tracks, (H) *Uromastyx aegyptia* scat.

to only 82 known prior to our survey. The number of species recorded in the reserve is expected to increase as taxonomic revisions are completed and as further surveys are conducted. The information generated here will aid the conservation efforts of the reserve, provide

a baseline for long-term monitoring, and inform further regional research and conservation assessments.

The 1,551 records generated for the reserve add a substantial number of observations not only for KARNR, but also for the whole of the Arabian Peninsula, with an




Fig. 6. (A) *Bunopus tuberculatus*, (B) *Cyrtopodion scabrum*, (C) *Hemidactylus granosus*, (D) *Stenodactylus doriae*, (E) *Stenodactylus slevini*, (F) *Trigonodactylus arabicus*, (G) *Tropicolotes wolfgangboehmei*, (H) *Ptyodactylus cf. hasselquistii*.

8.59% increase based on 18,053 records for the region as of 2021 (Šmíd et al. 2021). Having a baseline of reptile species diversity (richness and abundance) is important from a conservation perspective, as the data can be used to monitor changes over time (Maritz et al. 2016). Without such diversity metrics, the decay of richness

and abundance, potentially signalling local extinction events (Cressey et al. 2015; Zipkin et al. 2020), will go undetected and the “shifting baseline” syndrome is likely to take effect (Pauly 1995; Soga and Gaston 2018). For KARNR, this established baseline places conservation managers and stakeholders in a position to monitor future




Fig. 7. (A) *Acanthodactylus boskianus*, (B) *Acanthodactylus hardyi*, (C) *Acanthodactylus ophiodurus*, (D) *Acanthodactylus schmidtii*, (E) *Mesalina brevirostris*, (F) *Mesalina* lineage 4, (G) *Mesalina saudi-arabica*, (H) *Diplometopon zarudnyi*.

trends in the reserve. In addition, this survey provides a benchmark for comparisons to other reserves in the Arabian Peninsula.

The KARNR lies in an area predicted to have high reptile species richness of between 20 to 40 species (Alatawi et al. 2020; Šmíd et al. 2021). The 34 species

confirmed to occur in the reserve are within these bounds of the richness projection, but more species may potentially occur here, particularly in the southern areas (Alatawi et al. 2020; Šmíd et al. 2021). Species richness was not evenly distributed over the different broad habitat types. Hills, outcrops, and escarpments had the highest




Fig. 8. (A) *Chalcides ocellatus*, (B) *Scincus mitranus*, (C) *Scincus conirostris*, (D) *Scincus conirostris* tracks, (E) *Varanus griseus*.

diversity in terms of species richness, followed by Wadis and rawdhats, and Plains. Dunes and dune valleys had the lowest species richness. This trend can be attributed to the habitat heterogeneity hypothesis, which states that regions with higher habitat heterogeneity have higher species diversity (MacArthur and MacArthur 1961).

Additional surveys within the reserve are likely to yield new species, but several of the species encountered during the surveys are also undergoing taxonomic revision. For example, tentative molecular analyses indicate that more than one species of the *Ptyodactylus hasselquistii* complex may occur within the KARNR,




Fig. 9. (A) *Eryx jayakari*, (B) *Eryx jayakari*, (C) *Spalerosophis diadema*, (D) *Malpolon moilensis*, (E) *Psammophis schokari*, (F) *Cerastes gasperettii*, (G) *Cerastes gasperettii*, (H) *Echis coloratus*.

thus the reserve's species list will need to be updated as these new taxa are formally described. Likewise, the *Mesalina guttulata* species complex (recorded during our survey as *Mesalina* lineage 4 following Sindaco et al. 2018) is also undergoing taxonomic revision.

The four species not previously recorded from the

KARNR region are *Trapelus agnetae*, *Hemidactylus granosus*, *Mesalina saudiarabica*, and *Diplometopon zarudnyi*. The occurrence of *M. saudiarabica* was confirmed by DNA barcoding, with three samples of this species occurring in the Ad Dahna sand stretch of the reserve. The occurrence of this species was unexpected

Reptiles of the King Abdulaziz Royal Nature Reserve




Fig. 10. Predictive species accumulation curve generated with the R package iNEXT. The solid line represents observations made during the 2022/23 survey period and the stippled line represents the extrapolation of the data to double the survey effort.

based on its previously known distribution that was restricted to the western deserts of Saudi Arabia (Šmid et al. 2017). Not only do these records represent a significant range extension of more than 600 km to the north-east from the previously known range, but they also indicate the need for more detailed taxonomic investigations and herpetofaunal surveys in this understudied region as well as in other parts of central Saudi Arabia that remain largely under-surveyed.

One of the most frequently detected species within the KARNR is *Uromastix aegyptia*. This species is widespread throughout the reserve and occurs in all broad habitat types. *Uromastix aegyptia* is listed as Vulnerable by the IUCN, with threats relating to




Fig. 11. Predictive species accumulation curves generated with the R package iNEXT for four different habitat types (Dunes and dune valleys; Hills, outcrops, and escarpments; Plains; and Wadis, rawdhats, and pans). The solid lines represent observations made during the surveys and the stippled lines represent the extrapolation of the data to double the survey effort.

increasing anthropogenic pressures such as agricultural expansion, overgrazing, and poaching (IUCN 2023). Since the reserve was established, proactive measures have been taken to prevent habitat degradation, and hunting of *U. aegyptia* within the reserve borders has been made illegal. This highlights the role of KARNR as an important conservation area for reptiles in general, but especially for this Threatened species. Climate change is thought to be an additional threat to *U. aegyptia*, with ecological niche modelling projections indicating that habitat suitability across the Arabian Peninsula will decrease by 2070 (Kechnebbou et al. 2021). Under the worst-case scenario, only extremely limited areas within the reserve are predicted to remain suitable for *U.*

Table 2. Summary of species diversity calculations performed using the species accumulation curves generated with the R package iNEXT. For the KARNR as a whole and each of the broadscale habitat types, the diversity indices (Species richness, Shannon, and Simpson) are summarized in terms observed and estimated diversity with the standard error and upper and lower confidence levels. The iNEXT completeness percentage was calculated based on the observed versus estimated diversity.

Assemblage	Diversity	Observed	Estimator	Standard error	LCL	UCL	iNEXT completeness
KARNR	Species richness	31.00	31.67	3.15	31.00	37.84	97.88
	Shannon diversity	8.39	8.48	0.29	7.92	9.04	98.94
	Simpson diversity	4.75	4.76	0.17	4.43	5.09	99.79
Dunes and dune valleys	Species richness	14.00	21.96	6.29	14.00	34.29	63.75
	Shannon diversity	8.17	8.60	0.59	7.45	9.75	94.99
	Simpson diversity	6.30	6.47	0.54	5.40	7.53	97.42
Hills, outcrops, and escarpments	Species richness	24.00	24.25	5.97	24.00	35.95	98.97
	Shannon diversity	9.68	10.07	0.65	8.80	11.34	96.11
	Simpson diversity	6.33	6.44	0.50	5.46	7.42	98.29
Plains	Species richness	21.00	23.66	7.34	21.00	38.05	88.75
	Shannon diversity	4.89	4.99	0.31	4.39	5.60	97.94
	Simpson diversity	3.39	3.40	0.14	3.13	3.68	99.60
Wadis, rawdhats, and pans	Species richness	21.00	22.12	4.35	21.00	30.65	94.93
	Shannon diversity	6.37	6.54	0.44	5.69	7.40	97.41
	Simpson diversity	3.37	3.39	0.25	2.90	3.88	99.45

aegyptia, while suitability remains across KARNR for the best-case scenario (Kechnebbou et al. 2021). As such, the KARNR is an important sanctuary for the long-term protection of this species. However, future conservation plans for the reserve should focus not only on mitigating the impacts of anthropogenic pressures, but also on developing a monitoring program to identify potential range reductions within the reserve borders in the context of climate change.

The other reptile of conservation concern found within the reserve is *Tropicolotes wolfgangboehmei*, which is listed as Data Deficient by the IUCN and considered one of the world's most range-restricted reptiles (IUCN 2023; Meiri et al. 2018). This species was previously known from only three localities (Šmíd et al. 2021). We observed two individuals of this species during our survey, one at a previously known site within the 10 km buffer (300 m from the reserve boundary), while the other represents a new locality record (25 km north-east from the known site). Both records are from the Hills, outcrops, and escarpments habitat type. Future surveys in the KARNR should target this range-restricted species at additional sites and investigate its ecological requirements to assess the extent of its protection within the reserve.

In conclusion, our herpetological survey in the KARNR contributes significantly to our knowledge of the reptile diversity in east-central Saudi Arabia and provides a robust baseline for the region. The occurrences of 34 reptile species have been confirmed for the reserve, including four not previously recorded. The survey effort assessment, through predictive species accumulation curves and diversity indices, indicates that our methodology was adequate by documenting an estimated 97.89% of reptile species present in the KARNR. However, we acknowledge that further field and genetic investigations may reveal additional species.

The distribution of species richness across different broad habitat types reveals insights into the importance of habitat heterogeneity in supporting reptile diversity. The KARNR is a protected area of significant conservation importance, both at the regional and global scales. Its protection is beneficial for species of conservation concern such as *Uromastix aegyptia* (VU) and *Tropicolotes wolfgangboehmei* (DD), and for the more common and widespread reptile species and assemblages in the region. Ongoing taxonomic revisions and the potential impacts of climate change on sensitive species emphasize the need for more rigorous monitoring and for the implementation of adaptive conservation strategies. Our findings not only contribute significantly to the herpetofaunal knowledge of the Arabian Peninsula but also serve as a foundational resource for conservation planning and assessments, long-term monitoring, and regional comparative studies.

Acknowledgments.—We are grateful to the King Abdulaziz Royal Nature Reserve for commissioning

this study and Namariq Engineering Services for their logistical support. All work was conducted under permit 23-43WP, issued by the Reserve Authority. A special thank you goes to Sayd Haj Aissa and Saeed Nezar Alam for their invaluable assistance in ensuring smooth logistics, local arrangements, and on-the-ground support during our survey. We extend our appreciation to Lukáš Pola, Doubravka Velenská, Denis Hlaváč, Marek Uvizl, and Vojtěch Waldhauser for their contributions to the lab work associated with DNA barcoding. JŠ acknowledges the support of the Czech Science Foundation (GAČR) under grant number 22-12757S. We also express our gratitude to Awatef Abiadh, Ben Urban, Caroline Vasicek Gaugris, Colleen Lindberg, Graeme Wolfaard, Lukas Niemand, Marco Alexandre, Nicole Burri, Petrus Rossouw, Retief Grobler, Rio Button, Saeed Nezar Alam, and Tobi van Loggerenberg for providing additional records while in the field. We thank the anonymous reviewers for their comments and suggestions.

Literature Cited

- Alatawi AS, Gilbert F, Reader T. 2020. Modelling terrestrial reptile species richness, distributions, and habitat suitability in Saudi Arabia. *Journal of Arid Environments* 178: 104153.
- Almalki KA, Al Mosallam MS, Aldajani TZ, Al-Namazi AA. 2022. Landform characterization of Saudi Arabia: towards a geomorphological map. *International Journal of Applied Earth Observation and Geoinformation* 112: 102945.
- Aloufi AA, Amr ZS. 2015. On the herpetofauna of the Province of Tabuk, northwest Saudi Arabia. *Herpetozoa* 27(3/4): 147–158.
- Aloufi AA, Amr ZS, Abu Baker MA. 2021. Reptiles and amphibians of Al Madinah Al Munawwarah Province, Saudi Arabia. *Russian Journal of Herpetology* 28(3): 123–137.
- Aloufi AA, Amr ZS, Baker MA. 2022. Reptiles from ‘Uruq Bani Ma’arid and Harat al Harrah protected areas in Saudi Arabia: reptiles from two protected areas in Saudi Arabia. *Herpetology Notes* 15: 483–491.
- Aloufi AA, Amr ZS, Abu Baker MA, Hamidan N. 2019. Diversity and conservation of terrestrial, freshwater, and marine reptiles and amphibians in Saudi Arabia. *Amphibian & Reptile Conservation* 13(2) [General Section]: 181–202 (e204).
- Al-Sadoon MK. 1988. Survey of the reptilian fauna of the Kingdom of Saudi Arabia. II. The lizard and amphisbaenian fauna of Riyadh Province. *Bulletin of the Maryland Herpetological Society* 24(3): 58–76.
- Al-Sadoon MK, Paray BA, Al-Otaibi HS. 2016. Survey of the reptilian fauna of the Kingdom of Saudi Arabia. V. The lizard fauna of Turaif region. *Saudi Journal of Biological Sciences* 23(5): 642–648.
- Al-Sadoon MK. 1989. Survey of the reptilian fauna of

- the Kingdom of Saudi Arabia. I. The snake fauna of the central region. *Journal of King Saud University - Science* 1(1): 2.
- Al-Sadoon MK, Paray BA, Al-Otaibi H. 2017. Survey of the reptilian fauna of the Kingdom of Saudi Arabia. VI. The snake fauna of Turaif region. *Saudi Journal of Biological Sciences* 24(4): 925–928.
- Alshammari AM, Busais SM, Ibrahim AA. 2017. Snakes in the Province of Ha'il, Kingdom of Saudi Arabia, including two new records. *Herpetozoa* 30(1–2): 59–63.
- Alshammari AM, Busais SM. 2020. Distribution of snakes in Ha'il Province, Saudi Arabia, with an identification key to the species. *Russian Journal of Herpetology* 27(1): 5–10.
- Alshammari AM, Ibrahim AA. 2015. Lizards and snakes in the historical Faid protected area (Faid Hema), Ha'il region, Saudi Arabia. *Herpetological Conservation and Biology* 10(3): 1,021–1,029.
- Carranza S, Els J, Burriel-Carranza B. 2021. *A Field Guide to the Reptiles of Oman*. Bloomsbury Publishing, New York, New York, USA. 226 p.
- Chao A, Gotelli NJ, Hsieh TC, Sander EL, Ma KH, Colwell RK, Ellison AM. 2014. Rarefaction and extrapolation with Hill numbers: a framework for sampling and estimation in species diversity studies. *Ecological Monographs* 84(1): 45–67.
- Chao A, Jost L. 2012. Coverage-based rarefaction and extrapolation: standardizing samples by completeness rather than size. *Ecology* 1(12): 2,533–2,547.
- Cox NA, Mallon D, Bowles P, Els J, Tognelli MF. 2012. *The Conservation Status and Distribution of Reptiles of the Arabian Peninsula*. IUCN, Cambridge, United Kingdom and Gland, Switzerland, and Environment and Protected Areas Authority, Sharjah, United Arab Emirates. 49 p.
- Cressey ER, Measey GJ, Tolley KA. 2015. Fading out of view: the enigmatic decline of Rose's Mountain Toad *Capensibufo rosei*. *Oryx* 49(3): 521–528.
- Cunningham PL. 2010. Checklist of terrestrial reptiles in three protected areas in the Kingdom of Saudi Arabia. *Herpetological Review* 41(1): 25–28.
- Farag AA, Banaja AA. 1980. Amphibians and reptiles from the western region of Saudi Arabia. *Bulletin of Science, King Abdulaziz University* 4: 5–29.
- Ficetola GF, Bonardi A, Sindaco R, Padoa-Schioppa E. 2013. Estimating patterns of reptile biodiversity in remote regions. *Journal of Biogeography* 40(6): 1,202–1,211.
- Gasperetti J. 1988. Snakes of Arabia. *Fauna of Saudi Arabia* 9: 169–450.
- Ghazanfar SA, Fisher M, Editors. 1998. *Vegetation of the Arabian Peninsula (Number 25)*. Springer Science & Business Media, Berlin, Germany. 373 p.
- Hsieh TC, Ma KH, Chao A. 2016. iNEXT: an R package for rarefaction and extrapolation of species diversity (Hill numbers). *Methods in Ecology and Evolution* 7(12): 1,451–1,456.
- International Union of Conservation for Nature (IUCN). 2023. IUCN Red List of Threatened Species. IUCN: Gland, Switzerland. Available: <https://www.iucnredlist.org/> [Accessed: 7 July 2023].
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, et al. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28(12): 1,647–1,649.
- Kechnebbou M, de Carvalho DL, da Silva PH, Silva DP. 2021. Global warming drives range shifts in spiny-tailed lizards (Squamata: Agamidae: *Uromastyx*) in the African and Arabian deserts. *Journal of Arid Environments* 191: 104522.
- Kordges T. 1998. Die reptilian Fauna des Thumama Nature Park bei Riyadh, Saudi Arabien. *Faunistische Abhandlungen Staatliches Museum für Tierkunde Dresden* 21: 67–83.
- MacArthur RH, MacArthur JW. 1961. On bird species diversity. *Ecology* 42(3): 594–598.
- Margules CR, Pressey RL. 2000. Systematic conservation planning. *Nature* 405(6783): 243–253.
- Maritz B, Penner J, Martins M, Crnobrnja-Isailovic J, Spear S, Alencar LRV, Sigala-Rodriguez J, Messenger K, Clark RW, Soorae P, et al. 2016. Identifying global priorities for the conservation of vipers. *Biological Conservation* 204: 94–102.
- Meiri S, Bauer AM, Allison A, Castro-Herrera F, Chirio L, Colli G, Roll U. 2018. Extinct, obscure, or imaginary: the lizard species with the smallest ranges. *Diversity and Distributions* 24(2): 262–273.
- Mendes DM, de Freitas Leão R, Toledo LF. 2015. Drift fences in traps: theoretical evidence of effectiveness of the two most common arrays applied to terrestrial tetrapods. *Natureza & Conservação* 13(1): 60–66.
- Mihoub JB, Henle K, Titeux N, Brotons L, Brummitt NA, Schmeller DS. 2017. Setting temporal baselines for biodiversity: the limits of available monitoring data for capturing the full impact of anthropogenic pressures. *Scientific Reports* 7(1): 41591.
- Moreno CE, Halffter G. 2001. On the measure of sampling effort used in species accumulation curves. *Journal of Applied Ecology* 38: 487–490.
- Peel MC, Finlayson BL, McMahon TA. 2007. Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences* 11(5): 1,633–1,644.
- QGIS Development Team. 2023. QGIS Geographic Information System. Open-Source Geospatial Foundation Project. Available: <http://qgis.osgeo.org> [Accessed: 17 March 2022].
- R Core Team. 2022. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available: <https://www.R-project.org/> [Accessed: 10 February 2022].

- Salvador A. 1982. *A Revision of the Lizards of the Genus Acanthodactylus* (Sauria: Lacertidae). *Bonner Zoologische Monographien, Nummer 16*. Zoologisches Forschungsinstitut und Museum Alexander Koenig, Bonn, Germany. 167 p.
- Sindaco R, Simo-Riudalbas M, Sacchi R, Carranza S. 2018. Systematics of the *Mesalina guttulata* species complex (Squamata: Lacertidae) from Arabia with the description of two new species. *Zootaxa* 4429(3): 513–547.
- Sinervo B, Mendez-De-La-Cruz F, Miles DB, Heulin B, Bastiaans E, Villagrán-Santa Cruz M, Lara-Resendiz R, Martínez-Méndez N, Calderón-Espinosa ML, Meza-Lázaro RN, et al. 2010. Erosion of lizard diversity by climate change and altered thermal niches. *Science* 328(5980): 894–899.
- Šmíd J, Moravec J, Gvoždík V, Štundl J, Frynta D, Lymberakis P, Kapli P, Wilms T, Schmitz A, Shobrak M, et al. 2017. Cutting the Gordian knot: phylogenetic and ecological diversification of the *Mesalina brevirostris* species complex (Squamata, Lacertidae). *Zoologica Scripta* 46(6): 649–664.
- Šmíd J, Sindaco R, Shobrak M, Busais S, Tamar K, Aghová T, Simó-Riudalbas M, Tarroso P, Geniez P, Crochet PA, et al. 2021. Diversity patterns and evolutionary history of Arabian squamates. *Journal of Biogeography* 48(5): 1,183–1,199.
- Soga M, Gaston KJ. 2018. Shifting baseline syndrome: causes, consequences, and implications. *Frontiers in Ecology and the Environment* 16(4): 222–230.
- Tamar K, Uvizl M, Shobrak M, Almutairi M, Busais S, Salim AFA, AlGethami RHM, AlGethami AR, Alanazi ASK, Alsubaie SD, et al. 2023. A new species of *Pseudotrapelus* (Reptilia: Squamata: Agamidae) from Central Arabia. *Vertebrate Zoology* 73: 1,033–1,045.
- Thompson GG, Thompson SA, Withers PC, Fraser J. 2007. Determining adequate trapping effort and species richness using species accumulation curves for environmental impact assessments. *Austral Ecology* 32(5): 570–580.
- Van Rooyen M, Van Rooyen N, Gaugris JY. 2023. *Study of the Ecological Aspects – Botanical Studies of the Master Plan Section of the King Abdulaziz Royal Nature Reserve*. FLORA FAUNA & MAN, Ecological Services Limited, Tortola, British Virgin Islands. 121 p.
- Vincent P. 2008. *Saudi Arabia: an Environmental Overview*. CRC Press, London, United Kingdom. 332 p.
- Zipkin EF, DiRenzo GV, Ray JM, Rossman S, Lips KR. 2020. Tropical snake diversity collapses after widespread amphibian loss. *Science* 367(6479): 814–816.



Ryan van Huyssteen is a herpetology consultant at FLORA FAUNA & MAN, Ecological Services Ltd., specializing in herpetofauna surveys with experience in Africa and the Arabian Peninsula.



Melissa Petford is a conservation biologist and ecologist at FLORA FAUNA & MAN, Ecological Services Ltd., specializing in herpetology. She earned her Ph.D. from the University of the Witwatersrand (Johannesburg, South Africa) in 2019, and completed a postdoctoral position at the South African National Biodiversity Institute in February 2023. Melissa’s primary research focuses on understanding the impacts of anthropogenic activities on reptile distribution and ecology.



Marius Burger is an associate consultant with FLORA FAUNA & MAN, Ecological Services Ltd., a member of the IUCN SSC Amphibian Specialist Group (Sub-Saharan Africa), a member of IUCN SSC Snake and Lizard Red List Authority, and an extraordinary lecturer with the African Amphibian Conservation Research Group at North-West University (Potchefstroom, South Africa). He conducts herpetofaunal surveys and environmental impact assessments in Africa and the Arabian Peninsula.



Jiří Šmíd is a herpetologist and phylogeneticist at the National Museum in Prague, Czech Republic. His main line of research focuses on the reptiles of the Arabian Peninsula and its surroundings. He has described many endemic species from this region, including geckos, lacertids, and agamids.

Reptiles of the King Abdulaziz Royal Nature Reserve



Phoebe Mottram has an M.Sc. degree in Conservation Biology from the University of Cape Town, South Africa, and worked as a mammalian biologist for the past four years, focusing on biodiversity baseline surveys for development projects across western and central Africa. During 2023, she worked in Saudi Arabia where she focused on biodiversity baseline surveys and protected area planning for the King Abdulaziz Royal Nature Reserve. She presently works on biodiversity footprinting strategy development for large businesses.



Abdulrahman S. Alzahrani is an environmental scientist and acting wildlife manager with over 10 years of experience. He holds a Bachelor's degree in Environmental Science and a Master's degree in Environmental Science and Technology from RMIT University in Melbourne, Australia. Throughout his academic and professional journey, Abdulrahman has been actively involved in environmental research, focusing on areas such as water and soil quality assessment, biodiversity assessment, land conservation, and pollution assessments. His Master's thesis examined the effectiveness of stormwater wetlands in pollution removal and their role in supporting wildlife habitat in Geelong, Victoria, Australia. Abdulrahman is a certified Global and Regional Red List Assessor by the International Union for Conservation of Nature (IUCN), which demonstrates his expertise in conservation biology and his ability to assess the conservation status of species and ecosystems. Abdulrahman has contributed to projects aimed at preserving biodiversity, mitigating pollution, and promoting sustainable land management practices. He collaborates with various stakeholders to develop innovative solutions for environmental challenges.



Abdullah M. Alowaifeer is an environmental scientist interested in conservation, environmental contaminants, environmental risk assessment, and environmental rehabilitation. While pursuing his Ph.D. in Ecology at Montana State University (Bozeman, Montana, USA), he was involved in various projects that led him to gain theoretical and technical expertise in experimental design, sampling, and sample analysis. Abdullah did his Ph.D. thesis project on elemental cycling in Yellowstone National Park in the USA. His work was published and recognized as one of the top 10 Exceptional Papers of 2023 in the journal *Environmental Toxicology and Chemistry*. Abdullah has published over 20 scientific papers on various topics related to his work interests. He is currently the General Manager of Environmental Sustainability at King Abdulaziz Royal Nature Reserve in Saudi Arabia.



Jerome Gaugris is an accredited conservation scientist, ecologist, and wildlife manager. He is committed to providing practical, financially viable, and socially acceptable solutions that integrate human activities compatible with the long-term goal of biodiversity and ecosystem services conservation. Jerome has over 20 years of international professional experience focused on designing and implementing more than 100 scientifically grounded studies for ecological investigations and the sustainable use of renewable natural resources. Jerome is a vetted expert on Biodiversity and Sustainable Land Management for the United Nations Development Programme (UNDP) Africa, Caribbean and Pacific (ACP), and a vetted expert on Reduced Emissions from Deforestation and Forest Degradation (REDD+) for the Central African Forestry Initiative (CAFI).